

A NUTRITIONAL SUPPLEMENT, ADDED TO THE DRINKING WATER OF LAYING HENS, THAT HELPS IMPROVE THE NATURAL RESISTANCE OF POULTRY THROUGH NATURAL PLANT EXTRACTS.



APPLICATION TO POULTRY DRINKING WATER



DILUTION 500mL / 1000L

Building 69, Third Avenue, Pensnett Trading Estate, Kingswinford, West Midlands DY6 7FD Tel: 01384 404242 Email: sales@lodi-uk.com www.lodi-uk.com

Batch Nr. / Exp. date :

GMP021417 Armosa BV NU-CHI-02007 PA-LAB-03235



COMPLEMENTARY FEED ADDITIVE FOR LAYERS

- Liquid product to administer to poultry through drinking water.
- Supports the natural resistance of the hens.
- Ideal for laying hens and promotes a healthy response.
- Nutrients support birds during times of stress or anaemia.
- Natural ingredients suitable for organic farming

Deferder Pouliny Tonic contributies to support the natural resistance of layers. The natural ingredients provide a positive impact to layer's own four weeks later. R defences and reinforce the bird's natural antibioties to promote a healthy or sneeufive days. during times of stress or ranemia. STANDARD C

Defender Poultry Tonic is a liquid feed additive to administer in drinking water. Ideally, introduce (Protocol 1) at the start of the new flock, but can introduce at any point (Protocol 2) during the life cycle. In take guaranteed thanks to the application in drinking water. Easy to apply.

PRODUCT INFORMATION: Composition per L: Magnesium etholder, Stodium chioride, Vitamin C, Plant extracts (Origanum vulgare, Thymus vulgaris, Inula helenium, Litsea cubéba). RECOMMENDATIONS FOR USE:

DEFENDER POULTRY TONIC

must be administered in drinking water at the dosage of 0,5 L for 1000 L of water, at the rate of 200 ml per hen.

Protocol 1. Continuous application from flock introduction:

- 1 day per week, for the first 6 weeks. Renew this treatment every month for 2 consecutive days.

Protocol 2. Intermediate application in case of deficiencies: - For 4 consecutive days plus 2 days three weeks later and 2 days

Four weeks later. Renew this treatment every 15 to 20 weeks for 4 consecutive days.

STANDARD DEFENDER POULTRY TONIC TREATMENT PROTOCOL:

An effective protocol should always be farm specific. Ask our advice in case of increased infection pressure.

STORAGE/DISPOSAL: Store in a cool and dry place, between 5**Orm4/25*°C, away from direct light and heat sources. Defender Poultry Tonic should not be stored near electrical devices.