

**RACAN
FORCE PASTE**

Brodifacoum
(CAS 56073-10-0): 0.0025%
Warning: H373 May cause damage
to organs (blood) through
prolonged or repeated exposure.

P102: Keep out of reach of children.
P103: Read label before use. P220:
Keep/store away from food, drink
and animal feedingstuffs. P260: Avoid
breathing dust. P262: Do not get on
skin. P270: Do not eat, drink or smoke
when using this product. P301+310:
IF SWALLOWED: Immediately call a
POISON CENTRE or doctor/physician.
P314: Get medical advice/attention if
you feel unwell. P404+405: Store locked up
in a closed container. P501: Dispose of
contents/container in accordance with
applicable regulations.

**LODI UK - Building 69, Third Avenue,
Pensnett Trading Estate, Kingswinford,
West Midlands DY6 7FD, UK**
sales@lodi-uk.com www.lodi-uk.com

ADVICE

Racan Force Paste sachets are highly palatable
and lethal to rats and mice. The active ingredient,
Brodifacoum, is the UK's strongest active for rat
and mouse control. Rodent infestations will be
controlled within 4-10 days. Please ensure sachets
are applied and fixed securely in the Racan Rat
and Mouse Killer Station (not supplied) and DO NOT
come into contact with Children or Pets.



**RACAN
FORCE PASTE**

LodiUK

RAT & MOUSE KILLER PASTEBAIT



**SINGLE FEED
ACTIVE TECHNOLOGY**

ONE FEED - LETHAL DOSE



150g

**15 x 10g
PASTE
SACHETS**



**RACAN
FORCE PASTE**



**SINGLE FEED
ACTIVE TECHNOLOGY
FOR RATS AND MICE**

Our pioneering **Single Feed Active Technology** is specifically designed by our professional team to ensure users have the fastest, most lethal rodent control bait on the market. Rodents hit their lethal dose from as little as one feed, resulting in the quickest possible death. This product contains the active 'brodifacoum'.

**This pack contains
15 x 10g paste sachets**
produced to be highly
palatable and lethal to rats
and mice.

RACAN FORCE PASTE:
UFI: TK2W-8230-T00U-DU5C



0 700461 822905 >

RACAN FORCE PASTE
Paste bait containing 0.0025% w/w of Brodifacoum

**RACAN
FORCE PASTE**

Authorisation No: GB-2016-1058-0001 and NI-2016-1058-0001
Authorisation holder: Lodi-UK Building 69, Third Avenue, Pensnett
Trading Estate, Kingswinford, West Midlands DY6 7FD
Lodi SAS Parc d'Activités des Quatre Routes, 35380 Grand-Fougeray,
France

User category: For general public use.

Field of use: Indoor use for mice and rats / outdoor around buildings
for rats only.

Application method: Ready-to-use bait to be used in tamper-resistant
bait stations

Tamper-resistant bait stations must be:

- Strong enough to prevent entry or destruction by dogs.

- Strong enough to prevent entry or destruction by children under six years
of age using hands, feet or objects reasonably expected to be available in
the use area (e.g. household objects, toys, sticks, stones etc)

- Lockable or sealable so that children and dogs cannot gain access
through the opening or mechanisms used to fill the bait compartments.

- Resistant to destruction or weakening from exposure to typical non-
catastrophic weather (e.g. direct sunlight, extremes of temperature and
humidity, rain, snow etc).

Target organisms: *Rattus norvegicus* (Brown Rat) ; *Rattus rattus* (Black or
Roof Rat) ; *Mus musculus* (House mouse).

Application rate(s) and frequency:

For rats: 60g of bait per bait station. If more than one bait station
is needed, the minimum distance between bait stations should be
5-10m apart. For mice: 10g of bait per bait station. If more than 1 bait
station is needed, the minimum distance between bait stations should
be 3-5m apart.

Use-specific instructions for use: Mice: The bait stations should be
visited at least every 2 to 3 days at the beginning of the treatment and at
least weekly afterwards, in order to check whether the bait is accepted,
the bait stations are intact and to remove rodent bodies. Re-fill bait when
necessary. Rats: Place the bait stations in areas non-labile to flooding.
Replace any bait in a bait station in which bait has been damaged by
water or contaminated by dirt. The bait stations should be visited only 5 to
7 days after the beginning of the treatment and at least weekly afterwards,
in order to check whether the bait is accepted, the bait stations are intact
and to remove rodent bodies. Re-fill bait when necessary.

Instructions for use: Read and follow the product information as well
as any information accompanying the product or provided at the point of
sale before using it. Prior to the use of rodenticide products, non-chemical
control methods (e.g. traps) should be considered. Remove food which
is readily attainable for rodents (e.g. spilled grain or food waste). Apart
from this, do not clean up the infested area just before the treatment, as
this only disturbs the rodent population and makes bait acceptance more
difficult to achieve. Bait stations should be placed in the immediate vicinity
where rodent activity has been observed (e.g. travel paths, nesting sites,
tearlots, holes, burrows etc.). Where possible, bait stations must be fixed
to the ground or other structures. Do not open the sachets containing
the bait. Place bait stations out of the reach of children, birds, pets, farm
animals and other nontarget animals. Place bait stations away from food,
drink and animal feeding stuffs, as well as from utensils or surfaces that
have contact with these. Do not place bait stations near water drainage
systems where they can come into contact with water. When using the

product do not eat, drink or smoke. Wash hands and directly exposed skin
after using the product. Remove the remaining bait or the bait stations at
the end of the treatment period.

Risk mitigation measures: Consider preventive control measures
(plug holes, remove potential food and drinking as far as possible) to
improve product intake and reduce the likelihood of reinfestation. Do not
use anticoagulant rodenticides as permanent baits (e.g. for prevention of
rodent infestation or to detect rodent activity). Using this product should
eliminate rodents within 35 days. In case of suspected lack of efficacy
by the end of the treatment (i.e. rodent activity is still observed), the user
should seek advice from the product supplier or call a pest control service.
Search for and remove dead rodents during treatment, at least as often as
baiting stations are inspected. Dispose dead rodents in accordance with
local requirements. In the UK, poisoned rodents should be double-bagged
using plastic bags and either disposed of in a household waste bin with a
secure lid to prevent access of wildlife or pets or collected by a specialist
waste contractor or the local authority.

**Particulars of likely direct or indirect effects, first aid instructions
and emergency measures to protect the environment:** This product
contains an anticoagulant substance. If ingested, symptoms, which may
be delayed, may include nosebleed and bleeding gums. In severe cases,
there may be bruising and blood present in the faeces or urine. Antidote:
Vitamin K1 administered by medical/veterinary personnel only. In case of:
Dermal exposure, wash skin with water and then with water and soap.
Eye exposure, rinse eyes with eyes-rinse liquid or water. Keep eyes lids
open at least 10 minutes. Oral exposure, rinse mouth carefully with water.
Never give anything by mouth to unconscious person. Do not provoke
vomiting. If swallowed, seek medical advice immediately and show the
product's container or label. UK medical professionals should contact the
National Poisons Information Service (www.npis.org) for further advice.
Contact a veterinary surgeon in case of ingestion by a pet. Bait stations
must be labelled with the following information: "do not move or open";
"contains a rodenticide"; "product name or authorisation number";
"active substances"; and "in case of accident, call a poison centre".
Hazardous to wildlife.

Instructions for safe disposal of the product and its packaging:
At the end of the treatment, dispose unopened bait and the packaging in
accordance with local requirements. In the UK waste bait should be double
bagged in plastic bags and disposed of in a household waste bin with a
secure lid to prevent access of wildlife or pets or taken to a civic amenity
site. For information on civic amenity sites contact the local authority.
**Conditions of storage and shelf-life of the product under normal
conditions of storage:** Store in a dry, cool and well ventilated place.
Keep the container closed and away from direct sunlight. Store in places
prevented from the access of children, birds, pets and farm animals.
Shelf life: 2-years.

Because of their delayed mode of action, anticoagulant rodenticides take
from 4 to 10 days to be effective after consumption of the bait. Rodents
can be disease carriers. Do not touch dead rodents with bare hands, use
gloves or use tools such as long as long as when disposing them. This product
contains a bittering agent and a dye.
Lot No. and Expiry Date: See pack.

LodiUK

Marketed by LODI UK Limited:
Building 69, Third Avenue, Pensnett Trading Estate, Kingswinford,
West Midlands DY6 7FD Tel: 01384 404242 Email: sales@lodi-uk.com
www.lodi-uk.com

e 150g